

NLO QCD+EW for V+jets

Marek Schönherr*

Physik-Institut, Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland

E-mail: marek.schoenherr@physik.uzh.ch

Stefan Kallweit

Institut für Physik & PRISMA Cluster of Excellence, Johannes Gutenberg Universität, 55099 Mainz, Germany

E-mail: kallweit@uni-mainz.de

Jonas M. Lindert

Physik-Institut, Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland

E-mail: lindert@physik.uzh.ch

Philipp Maierhöfer

Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, 79104 Freiburg, Germany

E-mail: philipp.maierhoefer@physik.uni-freiburg.de

Stefano Pozzorini

Physik-Institut, Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland

E-mail: pozzorin@physik.uzh.ch

In this contribution recent results regarding the NLO electroweak corrections for vector boson production in association with jets are presented. Besides discussing the phenomenology of the fixed-order results, their incorporation in existing NLO QCD parton shower matched and merged calculations, which can directly be used in experimental analyses, will be shown.

Fourth Annual Large Hadron Collider Physics

13-18 June 2016

Lund, Sweden

*Speaker.

1. Introduction

The production of electroweak gauge bosons, frequently accompanied by one or multiple jets, plays a key role in the physics programme of the Large Hadron Collider. When decaying into leptons, the comparatively clean final state offers unique opportunities to test the Standard Model at the highest precisions. Simultaneously, vector boson production in association with multiple jets represents a large and therefore important background to many new physics searches.

As predictions for vector boson plus jets at next-to leading order (NLO) in QCD [1, 2, 3, 4, 5, 6, 7] are widely available, and the calculation of $V + 1$ jet final states is even known at next-to-next-to leading order (NNLO) [8, 9, 10, 11, 12, 13], electroweak (EW) corrections become increasingly important. They are relevant both for precision observables and measurements in the TeV regime where large Sudakov logarithms can lead to sizeable reductions of cross sections [14, 15, 16, 17, 18, 19]. While specific calculations for low multiplicity processes were available for some time [20, 21, 22, 23, 24], the recent automation of the generation of one-loop scattering amplitudes delivered the flexibility to calculate NLO EW corrections for processes with multiple jets in the final state [25, 26, 27, 28]. Automated implementations of the pure virtual Sudakov corrections are also available [29].

2. Next-to-leading order electroweak corrections

Next-to leading order electroweak corrections to (off-shell) vector boson ($V = W, Z, \gamma$) plus multijet production exhibit a rich internal structure. Apart from the occurrence of loop diagrams with multiple (different) internal masses, interferences from diagrams of different order in the QCD and EW couplings, g_s and e , are encountered as soon as four external quarks are present in the process. Thus, QCD and EW corrections cannot solely be classified as corrections due to real and virtual parton or photon emissions, respectively, but have to be defined through a power counting of the couplings involved: given a Born process at $\mathcal{O}(\alpha_s^n \alpha^m)$, NLO QCD corrections are defined as corrections of $\mathcal{O}(\alpha_s^{n+1} \alpha^m)$ while NLO EW corrections are of $\mathcal{O}(\alpha_s^n \alpha^{m+1})$. Consequently, NLO EW real corrections also involve processes with no external photons but an emergent pair of quarks or a gluon.

To calculate the NLO QCD and EW corrections to $W + n$ jets ($n = 1, 2, 3$) we use the fully automated frameworks of MUNICH+OPENLOOPS and SHERPA+OPENLOOPS. Therein, OPENLOOPS [30, 31] provides the renormalised one-loop matrix elements while either MUNICH or SHERPA [32, 33] arrange for phase space integration and process management. While MUNICH uses OPENLOOPS matrix elements for the Born and real emission matrix elements as well as in its Catani-Seymour [34, 35] subtraction, SHERPA employs its own internal matrix element generators AMEGIC [36] and COMIX [37]. This duality of approaches is used to provide independent cross checks of the results. OPENLOOPS uses the COLLIER [38] tensor reduction library, employing CUTTOOLS [39] and ONELOOP [40] for re-evaluating unstable phase space points.

Fig. 1 presents the NLO QCD and NLO EW corrections on the transverse momentum of the W boson and the leading jet in events with at least one jet. The inclusive corrections are shown on the left. As can be seen, the NLO QCD corrections alone amount to an increase in the cross section of several hundred percent at large p_T [41]. This effect originates in the opening




Figure 1: NLO QCD and EW corrections to the W -boson and leading jet transverse momenta in $pp \rightarrow W^+ j$ at 13 TeV at the LHC. While both types of corrections are dominated by real radiation $pp \rightarrow W^+ jj$ topologies featuring back-to-back dijet topologies and subsequently atypically large K -factors in the inclusive case (left), vetoing such configurations (right) recovers the standard behaviour.

of an alternative channel of production of a W boson in association with two jets encountered in the real corrections: a hard and nearly back-to-back dijet system radiates a comparably soft W boson. As these configurations can be interpreted as a NLO real EW boson emission corrections to dijet production, it inherits its large initial cross section and dominates in the high- p_T region of the leading jet, but not of the W boson. Consequently, its accuracy is reduced to be effectively leading order. Similar observations can be made for the NLO EW corrections. This artefact is best remedied by a multijet merged approach, treating the troublesome dijet plus W configurations at next-to leading order as well. In the absence of such a merging in a fixed-order calculation the offending contributions can simply be vetoed, e.g. by requiring the azimuthal separation of the two leading jets to be at most $\frac{3\pi}{4}$. The results for such a vetoed calculation are shown on the right hand side of Fig. 1. Here, modest NLO QCD and typical Sudakov-type NLO EW corrections are




Figure 2: NLO QCD and EW corrections to the W -boson and subleading and third leading jet transverse momenta, respectively, in $pp \rightarrow W^+ jj$ (left) and $pp \rightarrow W^+ jjj$ (right) at 13 TeV at the LHC.

recovered.

NLO QCD and EW corrections for the transverse momentum and the second and third leading jet in the higher multiplicity $W + 2j$ and $W + 3j$ events, respectively, are displayed in Fig. 2. As the Sudakov-type NLO EW corrections in the TeV regime become smaller for each subleading jet, so they do for the W boson itself. The results of [26] using on-shell bosons were quantitatively reproduced with an off-shell calculation using the full leptonic final state in [27].

These results were then directly used to produce theoretical predictions for the Z/γ -ratio in dependence on the (reconstructed) boson transverse momentum [44, 42, 45] in events with at least one jet accompanying the reconstructed Z boson or the photon, cf. Fig. 3 (left). As can be seen, the pure NLO QCD calculation predicts a slightly too large ratio. The electroweak correction, acting differently in both processes due to the different EW charge of the Z boson as compared to the photon, lower the ratio and improve the data description significantly.

Similarly, analyses focussed on finding W boson emissions off high- p_T jets [46], simplified




Figure 3: *Left:* Ratio of the transverse momentum of the reconstructed Z-boson and the photon in $pp \rightarrow \ell^+ \ell^- j$ and $pp \rightarrow \gamma j$, respectively, compared to data taken by the CMS collaboration at the LHC at 8 TeV [42]. *Right:* Angular separation of the muon and the closest jet in $pp \rightarrow W + jets$ production with $p_T^{j_1} > 500$ GeV at the LHC at 8 TeV, a comparison to data can be found in [43].

for the limited reach at 8 TeV, primarily measure the angular separation of the muon and the closest jet. As in this observables the region $\Delta R \gtrsim \pi$ is already defined in $W + 1j$ production but $\Delta R \lesssim \pi$ is only populated with $W + 2j$ events, an exclusive-sums approach (using $p_T^{j_2, \text{cut}} = 100$ GeV as discriminator) is taken. Fig. 3 (right) displays the NLO QCD+EW prediction for this observable including comparatively large subleading Born contributions. The yet unpublished study in [43] shows excellent agreement with data, especially where electroweak corrections are sizeable.

3. Electroweak corrections in particle-level event generation

The last section demonstrated the size and impact of next-to leading order electroweak corrections. However, they need to be incorporated in particle level event generators in order to be directly applied in experimental analyses. As a proper generic NLO QCD+EW matching to parton showers is still to be formulated, the following outlines an approximation that is designed to capture the dominant Sudakov-type corrections in the TeV regime as well as non-logarithmic ones over the whole phase space.

The basis for this approximation is SHERPA's MEPS@NLO [47, 48, 49, 50] multijet merging algorithm at NLO QCD accuracy. As it bases on an MC@NLO-type approach [51, 52, 53, 54, 55, 56] for matching the NLO QCD computation to the parton shower [57], its \bar{B} -function, labelled \bar{B}_{QCD} in the following, can be extended for the n parton final state to

$$\bar{B}_{n, \text{QCD+EW}_{\text{virt}}}(\Phi_n) = \bar{B}_{n, \text{QCD}}(\Phi_n) + V_{n, \text{EW}}(\Phi_n) + I_{n, \text{EW}}(\Phi_n) + B_{n, \text{mix}}(\Phi_n). \quad (3.1)$$

Herein, $V_{n, \text{EW}}(\Phi_n)$ are the renormalised one-loop electroweak corrections while $I_{n, \text{EW}}(\Phi_n)$ are the approximated real emission corrections integrated over the one-particle emission phase space,




Figure 4: Incorporation of approximate NLO EW corrections into the standard NLO QCD multijet merging for the transverse momenta of the reconstructed W -boson (left) and leading jet (right) in $pp \rightarrow \ell^- \bar{\nu}_\ell + jets$ at 13 TeV at the LHC.




Figure 5: Dependence of the reconstructed W -boson (left) and leading jet (right) transverse momenta on the merging parameter Q_{cut} in the LO (bottom) and NLO (top) QCD multijet merged calculation for $pp \rightarrow \ell^- \bar{\nu}_\ell + jets$ at 13 TeV at the LHC.

such that their sum yields a finite result. Optionally, subleading Born contributions, $B_{n,\text{mix}}$, can be included when numerically relevant. This approach neglects the kinematical impact of real photon radiation dominant in e.g. leptonic observables. It can be recovered, however, by making use of

standard QED radiation tools [58] which have been shown to be superior to a fixed order calculation for these observables [45].

Fig. 4 now again shows the transverse momentum distribution of the W boson on the left hand side and the leading jet on the right hand side obtained when calculating W + jets production with the above outlined method including up to two jets at (approximate) next-to leading order accuracy. As can be seen, very similar quantitative electroweak corrections are recovered in the TeV regime. Further, due to the consistent treatment of one and two jet final states the issue of potentially large corrections at large transverse momenta has been resolved in the merged calculation and the typical behaviour for the leading jet is recovered. For this observable, also the subleading Born contributions are of importance as they significantly reduce the impact of the electroweak corrections due to their opposite sign.

For the above prediction the standard merging cut of $Q_{\text{cut}} = 20 \text{ GeV}$ has been used. Following the Q_{cut} -dependence analysis of the method provided in [49, 50] large uncanceled logarithms could in principle emerge. A direct quantification through varying the merging parameter Q_{cut} yields the results presented in Fig. 5, showing negligible dependences for a large range of Q_{cut} values if varied around a sufficiently small central value.

4. Conclusions

The automation of NLO EW corrections within the MUNICH+OPENLOOPS and SHERPA+OPENLOOPS frameworks provide indispensable input for many high precision or TeV regime analyses at Run II of the LHC. A method for incorporation of approximate NLO EW corrections into established multijet merging methods at NLO QCD accuracy has been introduced and is publicly available within SHERPA-2.2.1. This formulation allows for particle level, and therefore potentially detector simulated, fully realistic event simulation. It represents a first, albeit very useful, step towards a complete NLO QCD+EW matching and multijet merging.

Acknowledgements. This research was supported in part by the Swiss National Science Foundation (SNF) under contract PP00P2-128552.

References

- [1] P. B. Arnold and M. H. Reno, *The Complete Computation of High $p(t)$ W and Z Production in 2nd Order QCD*, Nucl.Phys. **B319** (1989), 37.
- [2] P. B. Arnold, R. K. Ellis and M. Reno, *High $p(t)$ W and Z Production at the Tevatron*, Phys.Rev. **D40** (1989), 912.
- [3] J. M. Campbell and R. K. Ellis, *Next-to-leading order corrections to W^+ 2 jet and Z^+ 2 jet production at hadron colliders*, Phys.Rev. **D65** (2002), 113007, [[arXiv:hep-ph/0202176](#) [hep-ph]].
- [4] C. Berger, Z. Bern, L. J. Dixon et al., *Precise Predictions for W + 3 Jet Production at Hadron Colliders*, Phys.Rev.Lett. **102** (2009), 222001, [[arXiv:0902.2760](#) [hep-ph]].
- [5] R. K. Ellis, K. Melnikov and G. Zanderighi, *Generalized unitarity at work: first NLO QCD results for hadronic W + 3 jet production*, JHEP **0904** (2009), 077, [[arXiv:0901.4101](#) [hep-ph]].
- [6] C. Berger, Z. Bern, L. J. Dixon et al., *Precise Predictions for W + 4 Jet Production at the Large Hadron Collider*, Phys.Rev.Lett. **106** (2011), 092001, [[arXiv:1009.2338](#) [hep-ph]].

- [7] Z. Bern, L. Dixon, F. Febres Cordero, S. Höche, H. Ita et al., *Next-to-Leading Order $W + 5$ -Jet Production at the LHC*, Phys.Rev. **D88** (2013), no. 1, [014025](#), [[arXiv:1304.1253](#) [hep-ph]].
- [8] R. Boughezal, X. Liu and F. Petriello, *Phenomenology of the Z-boson plus jet process at NNLO*, [arXiv:1602.08140](#) [hep-ph].
- [9] R. Boughezal, X. Liu and F. Petriello, *W-boson plus jet differential distributions at NNLO in QCD*, [arXiv:1602.06965](#) [hep-ph].
- [10] R. Boughezal, X. Liu and F. Petriello, *A comparison of NNLO QCD predictions with 7 TeV ATLAS and CMS data for V+jet processes*, Phys. Lett. **B760** (2016), [6–13](#), [[arXiv:1602.05612](#) [hep-ph]].
- [11] R. Boughezal, C. Focke, X. Liu and F. Petriello, *W-boson production in association with a jet at next-to-next-to-leading order in perturbative QCD*, Phys. Rev. Lett. **115** (2015), no. 6, [062002](#), [[arXiv:1504.02131](#) [hep-ph]].
- [12] A. Gehrmann-De Ridder, T. Gehrmann, E. W. N. Glover, A. Huss and T. A. Morgan, *The NNLO QCD corrections to Z boson production at large transverse momentum*, JHEP **07** (2016), [133](#), [[arXiv:1605.04295](#) [hep-ph]].
- [13] A. Gehrmann-De Ridder, T. Gehrmann, E. W. N. Glover, A. Huss and T. A. Morgan, *Precise QCD predictions for the production of a Z boson in association with a hadronic jet*, Phys. Rev. Lett. **117** (2016), no. 2, [022001](#), [[arXiv:1507.02850](#) [hep-ph]].
- [14] V. S. Fadin, L. Lipatov, A. D. Martin and M. Melles, *Resummation of double logarithms in electroweak high-energy processes*, Phys.Rev. **D61** (2000), [094002](#), [[arXiv:hep-ph/9910338](#) [hep-ph]].
- [15] J. H. Kühn, A. Penin and V. A. Smirnov, *Summing up subleading Sudakov logarithms*, Eur.Phys.J. **C17** (2000), [97–105](#), [[arXiv:hep-ph/9912503](#) [hep-ph]].
- [16] A. Denner and S. Pozzorini, *One loop leading logarithms in electroweak radiative corrections. 1. Results*, Eur.Phys.J. **C18** (2001), [461–480](#), [[arXiv:hep-ph/0010201](#) [hep-ph]].
- [17] A. Denner and S. Pozzorini, *One loop leading logarithms in electroweak radiative corrections. 2. Factorization of collinear singularities*, Eur.Phys.J. **C21** (2001), [63–79](#), [[arXiv:hep-ph/0104127](#) [hep-ph]].
- [18] M. Ciafaloni, P. Ciafaloni and D. Comelli, *Bloch-Nordsieck violating electroweak corrections to inclusive TeV scale hard processes*, Phys.Rev.Lett. **84** (2000), [4810–4813](#), [[arXiv:hep-ph/0001142](#) [hep-ph]].
- [19] K. Mishra et al., *Electroweak Corrections at High Energies*, Proceedings, Community Summer Study 2013: Snowmass on the Mississippi (CSS2013): Minneapolis, MN, USA, July 29–August 6, 2013, 2013.
- [20] J. H. Kühn et al., *Electroweak corrections to hadronic photon production at large transverse momenta*, JHEP **03** (2006), [059](#), [[arXiv:hep-ph/0508253](#) [hep-ph]].
- [21] J. H. Kühn et al., *Electroweak corrections to hadronic production of W bosons at large transverse momenta*, Nucl.Phys. **B797** (2008), [27–77](#), [[arXiv:0708.0476](#) [hep-ph]].
- [22] A. Denner, S. Dittmaier, T. Kasprzik and A. Mück, *Electroweak corrections to W + jet hadroproduction including leptonic W-boson decays*, JHEP **0908** (2009), [075](#), [[arXiv:0906.1656](#) [hep-ph]].

- [23] A. Denner, S. Dittmaier, T. Kasprzik and A. Mück, *Electroweak corrections to dilepton + jet production at hadron colliders*, JHEP **06** (2011), 069, [[arXiv:1103.0914](#) [hep-ph]].
- [24] A. Denner, S. Dittmaier, T. Kasprzik and A. Mück, *Electroweak corrections to monojet production at the LHC*, Eur. Phys. J. **C73** (2013), no. 2, 2297, [[arXiv:1211.5078](#) [hep-ph]].
- [25] A. Denner, L. Hofer, A. Scharf and S. Uccirati, *Electroweak corrections to lepton pair production in association with two hard jets at the LHC*, JHEP **01** (2015), 094, [[arXiv:1411.0916](#) [hep-ph]].
- [26] S. Kallweit, J. M. Lindert, P. Maierhöfer, S. Pozzorini and M. Schönherr, *NLO electroweak automation and precise predictions for W+multijet production at the LHC*, JHEP **04** (2015), 012, [[arXiv:1412.5157](#) [hep-ph]].
- [27] S. Kallweit, J. M. Lindert, P. Maierhöfer, S. Pozzorini and M. Schönherr, *NLO QCD+EW predictions for V + jets including off-shell vector-boson decays and multijet merging*, JHEP **04** (2016), 021, [[arXiv:1511.08692](#) [hep-ph]].
- [28] M. Chiesa, N. Greiner and F. Tramontano, *Automation of electroweak corrections for LHC processes*, J. Phys. **G43** (2016), no. 1, 013002, [[arXiv:1507.08579](#) [hep-ph]].
- [29] M. Chiesa, G. Montagna, L. Barzè, M. Moretti, O. Nicrosini et al., *Electroweak Sudakov Corrections to New Physics Searches at the LHC*, Phys.Rev.Lett. **111** (2013), no. 12, 121801, [[arXiv:1305.6837](#) [hep-ph]].
- [30] F. Cascioli, P. Maierhöfer and S. Pozzorini, *Scattering Amplitudes with Open Loops*, Phys.Rev.Lett. **108** (2012), 111601, [[arXiv:1111.5206](#) [hep-ph]].
- [31] The OPENLOOPS one-loop generator by F. Cascioli, J. Lindert, P. Maierhöfer and S. Pozzorini is publicly available at <http://openloops.hepforge.org>.
- [32] T. Gleisberg and F. Krauss, *Automating dipole subtraction for QCD NLO calculations*, Eur.Phys.J. **C53** (2008), 501–523, [[arXiv:0709.2881](#) [hep-ph]].
- [33] T. Gleisberg et al., *Event generation with SHERPA 1.1*, JHEP **0902** (2009), 007, [[arXiv:0811.4622](#) [hep-ph]].
- [34] S. Catani and M. Seymour, *A General algorithm for calculating jet cross-sections in NLO QCD*, Nucl.Phys. **B485** (1997), 291–419, [[arXiv:hep-ph/9605323](#) [hep-ph]].
- [35] S. Catani et al., *The Dipole formalism for next-to-leading order QCD calculations with massive partons*, Nucl.Phys. **B627** (2002), 189–265, [[arXiv:hep-ph/0201036](#) [hep-ph]].
- [36] F. Krauss, R. Kuhn and G. Soff, *AMEGIC++ 1.0: A Matrix element generator in C++*, JHEP **0202** (2002), 044, [[arXiv:hep-ph/0109036](#) [hep-ph]].
- [37] T. Gleisberg and S. Höche, *Comix, a new matrix element generator*, JHEP **0812** (2008), 039, [[arXiv:0808.3674](#) [hep-ph]].
- [38] A. Denner, S. Dittmaier and L. Hofer, *Collier: a fortran-based Complex One-Loop Library in Extended Regularizations*, [arXiv:1604.06792](#) [hep-ph].
- [39] G. Ossola, C. G. Papadopoulos and R. Pittau, *CutTools: A Program implementing the OPP reduction method to compute one-loop amplitudes*, JHEP **0803** (2008), 042, [[arXiv:0711.3596](#) [hep-ph]].
- [40] A. van Hameren, *OneLOop: For the evaluation of one-loop scalar functions*, Comput.Phys.Commun. **182** (2011), 2427–2438, [[arXiv:1007.4716](#) [hep-ph]].
- [41] M. Rubin, G. P. Salam and S. Sapeta, *Giant QCD K-factors beyond NLO*, JHEP **09** (2010), 084, [[arXiv:1006.2144](#) [hep-ph]].

- [42] V. Khachatryan et al., CMS collaboration, *Comparison of the $Z/\gamma + \text{jets}$ to $\gamma + \text{jets}$ cross sections in pp collisions at $\sqrt{s} = 8 \text{ TeV}$* , JHEP **10** (2015), 128, [[arXiv:1505.06520](#) [hep-ex]], [Erratum: JHEP **04** (2016), 010].
- [43] M. Wu et al., *Observation and measurement of W emission collinear to high transverse momentum jets with the ATLAS detector*, ICHEP 2016 Chicago, 2016.
- [44] S. Kallweit, J. M. Lindert, S. Pozzorini, M. Schönherr and P. Maierhöfer, *NLO QCD+EW automation and precise predictions for V +multijet production*, Proceedings, 50th Rencontres de Moriond, QCD and high energy interactions: La Thuile, Italy, March 21–28, 2015, 2015, pp. 121–124.
- [45] J. R. Andersen et al., *Les Houches 2015: Physics at TeV Colliders Standard Model Working Group Report*, 9th Les Houches Workshop on Physics at TeV Colliders (PhysTeV 2015) Les Houches, France, June 1–19, 2015, 2016.
- [46] F. Krauss, P. Petrov, M. Schönherr and M. Spannowsky, *Measuring collinear W emissions inside jets*, Phys.Rev. **D89** (2014), 114006, [[arXiv:1403.4788](#) [hep-ph]].
- [47] S. Höche, F. Krauss, S. Schumann and F. Siegert, *QCD matrix elements and truncated showers*, JHEP **05** (2009), 053, [[arXiv:0903.1219](#) [hep-ph]].
- [48] S. Höche, F. Krauss, M. Schönherr and F. Siegert, *NLO matrix elements and truncated showers*, JHEP **08** (2011), 123, [[arXiv:1009.1127](#) [hep-ph]].
- [49] S. Höche, F. Krauss, M. Schönherr and F. Siegert, *QCD matrix elements + parton showers: The NLO case*, JHEP **1304** (2013), 027, [[arXiv:1207.5030](#) [hep-ph]].
- [50] T. Gehrmann, S. Höche, F. Krauss, M. Schönherr and F. Siegert, *NLO QCD matrix elements + parton showers in $e^+e^- \rightarrow \text{hadrons}$* , JHEP **01** (2013), 144, [[arXiv:1207.5031](#) [hep-ph]].
- [51] S. Frixione and B. R. Webber, *Matching NLO QCD computations and parton shower simulations*, JHEP **06** (2002), 029, [[hep-ph/0204244](#)].
- [52] S. Frixione, P. Nason and B. R. Webber, *Matching NLO QCD and parton showers in heavy flavour production*, JHEP **08** (2003), 007, [[hep-ph/0305252](#)].
- [53] S. Höche, F. Krauss, M. Schönherr and F. Siegert, *A critical appraisal of NLO+PS matching methods*, JHEP **09** (2012), 049, [[arXiv:1111.1220](#) [hep-ph]].
- [54] S. Höche, F. Krauss, M. Schönherr and F. Siegert, *W + n -jet predictions at the Large Hadron Collider at next-to-leading order matched with a parton shower*, Phys.Rev.Lett. **110** (2013), 052001, [[arXiv:1201.5882](#) [hep-ph]].
- [55] S. Höche and M. Schönherr, *Uncertainties in next-to-leading order plus parton shower matched simulations of inclusive jet and dijet production*, Phys.Rev. **D86** (2012), 094042, [[arXiv:1208.2815](#) [hep-ph]].
- [56] S. Höche, F. Krauss, M. Schönherr and F. Siegert, *Automating the POWHEG method in Sherpa*, JHEP **04** (2011), 024, [[arXiv:1008.5399](#) [hep-ph]].
- [57] S. Schumann and F. Krauss, *A parton shower algorithm based on Catani-Seymour dipole factorisation*, JHEP **03** (2008), 038, [[arXiv:0709.1027](#) [hep-ph]].
- [58] M. Schönherr and F. Krauss, *Soft Photon Radiation in Particle Decays in SHERPA*, JHEP **12** (2008), 018, [[arXiv:0810.5071](#) [hep-ph]].